



# Parent/Athlete Concussion Information Sheet

A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by bump, blow, or jolt to the head or body that causes the head and brain to move rapidly back and forth. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious.

## WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury.

If an athlete reports **one or more** symptoms of concussion listed below after a bump, blow, or jolt to

### Did You Know?

- Most concussions occur *without* loss of consciousness.
- Athletes who have, at any point in their lives, had a concussion have an increased risk for another concussion.
- Young children and teens are more likely to get a concussion and take longer to recover than adults.

the head or body, s/he should be kept out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it’s OK to return to play.

SIGNS OBSERVED BY COACHING STAFF	SYMPTOMS REPORTED BY ATHLETES
Appears dazed or stunned	Headache or “pressure” in head
Is confused about assignment or position	Nausea or vomiting
Forgets an instruction	Balance problems or dizziness
Is unsure of game, score, or opponent	Double or blurry vision
Moves clumsily	Sensitivity to light
Answers questions slowly	Sensitivity to noise
Loses consciousness ( <i>even briefly</i> )	Feeling sluggish, hazy, foggy, or groggy
Shows mood, behavior, or personality changes	Concentration or memory problems
Can’t recall events <i>prior</i> to hit or fall	Confusion
Can’t recall events <i>after</i> hit or fall	Just not “feeling right” or “feeling down”

## CONCUSSION DANGER SIGNS

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. An athlete should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that not only does not diminish, but gets worse
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless, or agitated
- Has unusual behavior
- Loses consciousness (*even a brief loss of consciousness should be taken seriously*)

## WHY SHOULD AN ATHLETE REPORT THEIR SYMPTOMS?

If an athlete has a concussion, his/her brain needs time to heal. While an athlete's brain is still healing, s/he is much more likely to have another concussion. Repeat concussions can increase the time it takes to recover. In rare cases, repeat concussions in young athletes can result in brain swelling or permanent damage to their brain. *They can even be fatal.*

### *Remember*

Concussions affect people differently. While most athletes with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

## WHAT SHOULD YOU DO IF YOU THINK YOUR ATHLETE HAS A CONCUSSION?

If you suspect that an athlete has a concussion, remove the athlete from play and seek medical attention. Do not try to judge the severity of the injury yourself. Keep the athlete out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.

Rest is key to helping an athlete recover from a concussion. Exercising or activities that involve a lot of concentration, such as studying, working on the computer, or playing video games, may cause concussion symptoms to reappear or get worse. After a concussion, returning to sports and school is a gradual process that should be carefully managed and monitored by a health care professional.

It's better to miss one game than the whole season. For more information on concussions, visit: [www.cdc.gov/Concussion](http://www.cdc.gov/Concussion).

\_\_\_\_\_  
Student-Athlete Name Printed

\_\_\_\_\_  
Student-Athlete Signature

\_\_\_\_\_  
Date

\_\_\_\_\_  
Parent or Legal Guardian Printed

\_\_\_\_\_  
Parent or Legal Guardian Signature

\_\_\_\_\_  
Date



# Sudden Cardiac Arrest

## Athlete/Parent/Guardian Sudden Cardiac Arrest Symptoms and Warning Signs Information Sheet and Acknowledgement of Receipt and Review Form

### What is sudden cardiac arrest?

Sudden cardiac arrest (SCA) is when the heart stops beating, suddenly and unexpectedly. When this happens, blood stops flowing to the brain and other vital organs. SCA doesn't just happen to adults; it takes the lives of students, too. However, the causes of sudden cardiac arrest in students and adults can be different. A youth athlete's SCA will likely result from an inherited condition, while an adult's SCA may be caused by either inherited or lifestyle issues. SCA is NOT a heart attack. A heart attack may cause SCA, but they are not the same. A heart attack is caused by a blockage that stops the flow of blood to the heart. SCA is a malfunction in the heart's electrical system, causing the heart to suddenly stop beating.

### How common is sudden cardiac arrest in the United States?

SCA is the #1 cause of death for adults in this country. There are about 300,000 cardiac arrests outside hospitals each year. About 2,000 patients under 25 die of SCA each year. It is the #1 cause of death for student athletes.

### Are there warning signs?

Although SCA happens unexpectedly, some people may have signs or symptoms, such as:

- fainting or seizures during exercise;
- unexplained shortness of breath;
- dizziness;
- extreme fatigue;
- chest pains; or
- racing heart.

These symptoms can be unclear in athletes, since people often confuse these warning signs with physical exhaustion. SCA can be prevented if the underlying causes can be diagnosed and treated.

### What are the risks of practicing or playing after experiencing these symptoms?

There are risks associated with continuing to practice or play after experiencing these symptoms. When the heart stops, so does the blood that flows to the brain and other vital organs. Death or permanent brain damage can occur in just a few minutes. Most people who experience SCA die from it.

**Public Chapter 325 – the Sudden Cardiac Arrest Prevention Act**

The act is intended to keep youth athletes safe while practicing or playing. The requirements of the act are:

- All youth athletes and their parents or guardians must read and sign this form. It must be returned to the school before participation in any athletic activity. A new form must be signed and returned each school year.
  
- The immediate removal of any youth athlete who passes out or faints while participating in an athletic activity, or who exhibits any of the following symptoms:
  - (i) Unexplained shortness of breath;
  - (ii) Chest pains;
  - (iii) Dizziness
  - (iv) Racing heart rate; or
  - (v) Extreme fatigue; and
  
- Establish as policy that a youth athlete who has been removed from play shall not return to the practice or competition during which the youth athlete experienced symptoms consistent with sudden cardiac arrest
  
- Before returning to practice or play in an athletic activity, the athlete must be evaluated by a Tennessee licensed medical doctor or an osteopathic physician. Clearance to full or graduated return to practice or play must be in writing.

*I have reviewed and understand the symptoms and warning signs of SCA.*

---

Signature of Student-Athlete

Print Student-Athlete's Name Date

---

Signature of Parent/Guardian

---

Print Parent/Guardian's Name Date